



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

RECORDS OF NORTHERN NEW JERSEY
FISHES.

Early in January, 1915, Prof. R. C. Osburn kindly forwarded me records of additions to the New York Aquarium obtained in New Jersey waters. These were derived from the notes of Mr. W. I. DeNyse. I am indebted to both these gentlemen for the opportunity of presenting the greater part of the following list: The localities are Belford (b), Sandy Hook (s), from off Seabright to off Long Branch (o) in 6 to 17 fathoms. Butler (bu), Boonton (bo), Lake Weequahic (w). I am also indebted to Mr. C. F. Silvester for notes on fishes at Elberon (e), Manasquan (m), North Long Branch (n), Long Branch (l) and Princeton in the canal (c), and Stony Brook (p). *Petromyzon marinus* (b, p.), *Carcharias taurus* (b), *Mustelus mustelus* (s, o, l.), *Eulamia milberti* (b, o.), *E. obscurus* (l), *Sphyrna zygaena* (l), *Squalus acanthias* (o, l, e, m.), *Raja erinacea* (b, o, l.), *R. ocellata* (l), *R. eglanteria* (o), *R. laevis*, (b, o, l.), *Narcobatus nobilianus* (s, one June, 1909), *Dasyatis centroura* (o), *Rhinoptera bonasus* (b, one October 13, 1912), *Acipenser sturio* (b), *Tarpan atlanticus* (b), *Dorosoma cepedianum* (c), *Pomolobus mediocris* (b), *P. pseudoharengus* (b), *Alosa sapidissima* (b), *Brevoortia tyrannus* (b), *Ameiurus nebulosus* (w), *Schilbeodes gyrimus* (p), *Abramis crysoleucas* (w, bu, bo.), *Notropis cornutus* (p), *Carassius auratus* (bu), *Catostomus commersonnii* (bo), *Erimyzon sucetta oblongus* (c, p.), *Esox americanus* (c), *E. reticulatus* (bu, Charlottesville), *Fundulus majalis* (s, Shrewsbury River), *F. heteroclitus macrolepidotus* (s), *F. diaphanus* (Bay-shore), *Cyprinodon variegatus* (s), *Tylosurus marinus* (s), *Hyporhamphus unifasciatus* (s), *Syngnathus fuscus* (m), *Hippocampus hudsonius* (l, s, b.), *Mugil curema* (b?, s.), *Scomber scombrus* (e), *Sarda sarda* (l, m.), *Thunnus thynnus* (l, n.), *Scomberomorus maculatus* (l), *Trichiurus lepturus* (l), *Istio-*

phorus nigricans (n), *Seriola zonata* (l, o.), *Pomatomus saltatrix* (s, b, n.), *Trachinotus falcatus* (b), *Rachycentron canadum* (b), *Poronotus triacanthus* (e), *Aphredoderus sayanus* (c), *Lepomis auritus* (c, p.), *Pomotis gibbosus* (bu, bo, w, p.), *Microp-terus dolomieu*, (w, bo, Millstone River below Rocky Hill), *M. salmoides* (bo), *Perca flavescens* (bu, bo, c, p.), *Boleosoma nigrum olmstedii* (p), *Roccus lineatus* (b, l, Barnegat), *Morone americana* (s), *Centropristis striatus* (s, b.), *Stenotomus chrysops* (s, b.), *Lagodon rhomboides* (b), *Mullus auratus* (s), *Cynoscion regalis* (l, s, b.), *Bairdiella chrysura* (s), *Leiostomus xanthurus* (s, b.), *Micropogon undulatus* (b), *Menticirrhus saxatilis* (s, e.), *Pogonias cromis* (b), *Tautogolabrus adspersus* (b), *Tautoga onitis* (l, s, b.), *Balistes carolinensis* (o), *Alutera schoepfii* (s, b, n.), *Stephanolepis hispidus* (b, s.), *Lagocephalus laevigatus* (o), *Spheroides maculatus* (o, s, b.), *Chilomycterus schoepfi* (s, b.), *Myoxocephalus octodecimspinosus* (n, m, o.), *M. aeneus* (s, b.), *Prionotus evolans strigatus* (s, b, o.), *P. carolinus* (o, s, b.), *Paralichthys dentatus* (b, s, o, l.), *Pseudopleuronectes americanus* (s), *Achirus fasciatus* (b), *Leptecheneis naucratis* (b), *Opsanus tau* (s, b, l.), *Astroscopus guttatus* (s), *Enchelyopus anguillaris* (o), *Rissola marginata* (b), *Gadus callarias* (o), *Pollachius virens* (o), *Microgadus tomcod* (s), *Urophycis regius* (s, l.), *U. chuss* (o), *U. tenuis* (o), *Merluccius bilinearis* (b, l.), *Lophius piscatorius* (o, l.), *Leptocephalus conger* (l).

HENRY W. FOWLER,
Philadelphia, Pa.

BURROWING HABITS OF THE BOX TURTLE.

Dr. Overton's note in COPEIA No. 26 on "Aquatic Habits of the Box Turtle" (*Cistudo carolina*) recalls to the writer observations on Long Island, which indicate periods of aestivation for this species.